

$\Delta + 300$ is a Bound on the Adjacent Vertex Distinguishing Edge Chromatic Number

Hamed Hatami

Department of Computer Science

University of Toronto

e-mail: hamed@cs.toronto.edu

Abstract

An adjacent vertex distinguishing edge-coloring or an *avd*-coloring of a simple graph G is a proper edge-coloring of G such that no pair of adjacent vertices meets the same set of colors. We prove that every graph with maximum degree Δ and with no isolated edges has an *avd*-coloring with at most $\Delta + 300$ colors, provided that $\Delta > 10^{20}$.

AMS SUBJECT CLASSIFICATION: 05C15

KEYWORDS: adjacent vertex distinguishing edge coloring, vertex distinguishing edge coloring, adjacent strong edge coloring, strong edge coloring.

1 Introduction

We follow [10] for terminologies and notations not defined here. For every vertex v in a graph G , $\deg_G(v)$ or $\deg(v)$ when there is no ambiguity denotes the degree of v in the graph G . For every partial edge coloring c of a graph G , and every vertex v of G , let $S_c(v)$ denote the set of the colors incident to v .

In a partial edge coloring c a vertex u is called **distinguishable**, if for every vertex v adjacent to u , $S_c(u) \neq S_c(v)$. An edge coloring c is called **adjacent vertex distinguishing** or an *avd*-coloring if every vertex is distinguishable. A *k-avd*-coloring is an *avd*-coloring using at most k colors. It is clear that every graph with isolated edges does not have any *avd*-coloring. The *avd*-chromatic number of a graph G , the minimum number of colors in an *avd*-coloring of G , is defined for every graph G without any isolated edge. Adjacent vertex distinguishing edge colorings are studied in [1, 2, 6, 11], where different names such as **adjacent strong edge coloring** [11] and **1-strong edge coloring** [1, 6] are used to refer to an *avd*-coloring. Adjacent vertex distinguishing edge colorings are related to vertex distinguishing edge colorings

in which the condition $S_c(u) \neq S_c(v)$ holds for every pair of vertices u and v , not necessarily adjacent. This concept has been studied in many papers, (see for example [3, 4, 5, 7]).

Another interesting problem arises when we drop the condition that the edge coloring is proper and allow the incident edges to have the same colors. The following theorem is proved in [8] by Karoński, Łuczak, and Thomason.

Theorem A. [8] *There exists a finite set of real numbers which can be used to weight the edges of any graph with no isolated edges so that adjacent vertices have different sums of incident edge weights.*

It is easy to observe that Theorem A is equivalent to the following theorem.

Theorem B. [restatement of Theorem A] *There is a finite set which can be used to color the edges of any graph with no isolated edges so that adjacent vertices meet different multisets (i.e. duplicate elements are allowed) of colors.*

This shows that by dropping the condition of being proper from the definition of *avd*-coloring, a constant number of colors would be sufficient. Obviously, when the edge colorings are required to be proper this is not the case. The following conjecture was made in [11].

Conjecture A. [11] *The avd-chromatic number of every simple connected graph G such that $G \neq C_5$ (the cycle of size 5) and $G \neq K_2$ is at most $\Delta + 2$.*

In [2] this conjecture is verified for bipartite graphs and graphs of maximum degree 3, and the following bound has been proved for general graphs.

Theorem C. [2] *If G is a graph with no isolated edges, then the avd-chromatic number of G is at most $\Delta + O(\log \chi(G))$.*

This was the best known bound so far. We asymptotically improve it by proving the following theorem.

Theorem 1 *If G is a graph with no isolated edges and maximum degree $\Delta > 10^{20}$, then the avd-chromatic number of G is at most $\Delta + 300$.*

We will use the probabilistic method to prove Theorem 1. The following tools of the probabilistic method will be used several times (see [9]).

Theorem D. [The Symmetric Lovász Local Lemma] *Let $A_1, \dots, A_n$ be events in a probability space, let for all i , $\Pr[A_i] = p$ and the event A_i is mutually independent of all events but at most d events. If $4pd < 1$, then $\Pr[\bigwedge \bar{A}_i] > 0$.*

Theorem E. [*The Chernoff Bound*] Suppose $BIN(n, p)$ is the sum of n independent Bernoulli variables each occurring with probability p . Then for any $0 < t \leq np$

$$\Pr[|BIN(n, p) - np| > t] < 2e^{-\frac{t^2}{3np}}.$$

Theorem F. [*Talagrand's Inequality*] Let X be a non-negative random variable, not identically 0, which is determined by n independent trials $T_1, \dots, T_n$, and satisfying the following for some $c, r > 0$

1. Changing the outcome of any one trial can affect X by at most c , and
2. For any s and any outcome of trials, if $X \geq s$, then there is a set of at most rs trials whose outcomes certify that $X \geq s$,

then for any $0 \leq t \leq \mathbf{E}[x]$,

$$\Pr[|X - \mathbf{E}[x]| > t + 60c\sqrt{r\mathbf{E}[x]}] \leq 4e^{-\frac{t^2}{8c^2r\mathbf{E}[X]}}.$$

Remark 1 Suppose that we have the upperbound $\mathbf{E}[X] \leq k$, and the conditions of Talagrand's Inequality hold for X . Then we might apply Talagrand's Inequality to the random variable $Y = X + k - \mathbf{E}[X]$, and obtain the inequality

$$\Pr[X > k + t + 60c\sqrt{rk}] \leq \Pr[Y > k + t + 60c\sqrt{rk}] \leq 4e^{-\frac{t^2}{8c^2rk}}.$$

Similar arguments hold for the Chernoff bound, and also for the cases that we know a lower bound for $\mathbf{E}[X]$ and want to apply Talagrand's Inequality or the Chernoff bound.

We will also use the following well known inequalities.

$$\left(\frac{a}{b}\right)^b \leq \binom{a}{b} = \binom{a}{a-b} \leq \left(\frac{ea}{b}\right)^b, \quad (1)$$

where a and b are natural numbers.

In Section 2 we present the proof of Theorem 1.

2 Proof of Theorem 1

The proof is probabilistic, and consists of three steps. The first step is similarly used in [6]. Let G be a graph with no isolated edges.

- In the first step we construct a loopless multigraph G' with multiplicity at most 2 so that $\Delta(G') = \Delta(G) = \Delta$, and if there exists a p -avd-coloring for G' , then G is also p -avd-colorable.

- Let H be the subgraph of G' induced by all vertices v such that $\deg(v) < \frac{\Delta}{3}$. In the second step we give a $(\Delta + 300)$ -edge coloring c of G' such that for every two adjacent vertices $u, v \in V(G') - V(H)$, $S_c(u) \neq S_c(v)$.
- In the third step we modify c by recoloring some edges in $E(H)$, and obtain a $(\Delta + 300)$ -*avd*-coloring, c_{final} of G' .

2.1 First Step

For every edge $e = uv$ in G such that $\deg(u), \deg(v) < \frac{\Delta}{3}$ and none of u and v has any other neighbor of degree less than $\frac{\Delta}{3}$, contract the edge e , i.e. remove e and identify u and v to a single vertex. Call this new multigraph G' . Obviously, $\Delta(G') = \Delta(G)$, and the multiplicity of G' is at most 2. Note that H , the subgraph of G' induced by all vertices v such that $\deg(v) < \frac{\Delta}{3}$, does not have any isolated edges.

Suppose that G' has a p -*avd*-coloring σ' . Every edge of G' has a corresponding edge in G . For every edge e of G' , color its corresponding edge in G by $\sigma'(e)$. Now all edges of G are colored except those edges which were contracted in the process of obtaining G' from G . For every such edge uv in G , since $\deg(v) + \deg(u) < \Delta$, there exist some available colors for uv . Color uv with one of these available colors. Trivially u and v meet different sets of colors and none of them has any other neighbor of the same degree. This implies that the obtained edge coloring of G is in fact an *avd*-coloring.

2.2 Second Step

An **unused edge** in a partial edge coloring c of a graph G is an edge which is not colored by c . We will refer to the graph U_c induced by all unused edges as the **unused graph** of c . The degree of a vertex in U_c is called its **unused degree**. We say that a color x is **available** for an unused edge uv , if x does not appear on any incident edge to uv .

As it is mentioned before, H is the subgraph of G' induced by all vertices v such that $\deg(v) < \frac{\Delta}{3}$. It is trivial that H has no multiple edges. We start with an arbitrary $\Delta + 2$ edge coloring σ of G' (which is guaranteed to exist by Vizing's Theorem), and then we apply a two-phased procedure to modify σ , and obtain an edge coloring c such that for every two adjacent vertices $u, v \in V(G') - V(H)$, $S_c(u) \neq S_c(v)$. The first phase is the following.

Phase I

1. Uncolor each edge $e \in E(G') - E(H)$ with probability $\frac{180}{\Delta}$.

2. For every vertex v which has unused degree greater than 290, recover the color of all unused edges which are incident to it. We say that v is **recovered**.

Let σ_1 be the partial coloring obtained after applying Phase **I**. We define the following sets.

- UC_v (uncolored) is the set of the edges which are incident to v and uncolored in Phase **I** (1). Note that these edges are not necessarily unused in σ_1 because their colors might be recovered in Phase **I** (2).
- R is the set of all recovered vertices.
- Q is the set of all vertices v such that $|UC_v| < 20$.
- T is the set of all vertices $v \in V(G') - V(H)$ such that there exists an edge $vw \in UC_v$ where w is recovered.
- L is the set of all vertices $v \in V(G') - V(H)$ such that $\deg_{U_{\sigma_1}}(v) < 20$, where U_{σ_1} is the unused graph of σ_1 .

Since we use randomness in Phase **I**, intuitively after applying this phase if a vertex has a large unused degree, then with a high probability it is distinguishable. However there are vertices in $V(G') - V(H)$ which have small unused degrees which we denote them by L . Later in Lemma 4 we will show that with a positive probability the vertices in L are rare and well-distributed in the graph. In fact we will prove that with positive probability:

- (a): For every vertex $v \in V(G') - V(H)$, we have $|N(v) \cap L| \leq \frac{\Delta}{100}$,
- (b): For every two adjacent vertices $u, v \in V(G') - V(H)$ such that $\deg(u) = \deg(v)$ and $u \notin L$, we have $|S_{\sigma_1}(u) \triangle S_{\sigma_1}(v)| \geq 10$, where for every two sets A and B , $A \triangle B = (A \setminus B) \cup (B \setminus A)$.

Note that L is a subset of the union of R , Q , and T . The following three lemmas provide the technical details needed to prove Lemma 4.

Lemma 1 *For every vertex $v \in V(G') - V(H)$, we have*

- (a) : $\Pr[v \in R] \leq \frac{1}{1000}$.
- (b) : $\Pr[v \in Q] \leq \frac{1}{1000}$.
- (c) : $\Pr[v \in T] \leq \frac{1}{1000}$.

Proof.

(a) Since $|UC_v|$ has Bernoulli distribution, knowing that $\frac{\Delta}{3} \leq \deg(v) \leq \Delta$, the Chernoff bound implies that

$$\Pr[v \in R] = \Pr[|UC_v| > 290] \leq 2e^{-\frac{(290-180)^2}{3 \times 180}} \leq \frac{1}{1000}.$$

(b) The worst case is when $\deg(v) = \frac{\Delta}{3}$. Then the Chernoff bound implies that

$$\Pr[v \in Q] = \Pr[|UC_v| < 20] \leq 2e^{-\frac{(180/3-20)^2}{180}} \leq \frac{1}{1000}.$$

(c) Consider an edge vw , where $v \in V(G') - V(H)$ and $w \in V(G')$. First we prove an upper bound for $\Pr[vw \in UC_v \wedge w \in R]$. The Chernoff bound implies that

$$\Pr[w \in R | vw \in UC_v] \leq 2e^{-\frac{(290-1-180)^2}{3 \times 180}} \leq \frac{1}{10^6}.$$

Hence

$$\Pr[vw \in UC_v \wedge w \in R] \leq \frac{1}{10^6} \frac{180}{\Delta} \leq \frac{1}{10^3 \Delta}.$$

Since v has at most Δ neighbors, we have

$$\Pr[v \in T] \leq \frac{1}{10^3 \Delta} \times \Delta \leq \frac{1}{1000}.$$

■

Lemma 2 For every vertex $v \in V(G') - V(H)$,

$$\Pr[|N(v) \cap L| > \Delta/100] \leq \frac{1}{\Delta^7}.$$

Proof. First notice that

$$|N(v) \cap L| \leq |N(v) \cap R| + |N(v) \cap T| + |N(v) \cap Q|.$$

Lemma 1 implies that

- $\mathbf{E}[|N(v) \cap R|] \leq \frac{\Delta}{1000},$
- $\mathbf{E}[|N(v) \cap Q|] \leq \frac{\Delta}{1000},$ and
- $\mathbf{E}[|N(v) \cap T|] \leq \frac{\Delta}{1000}.$

For edges $uv \in E(G') - E(H)$, consider the independent Bernoulli trials T_{uv} where the outcome of T_{uv} determines whether uv is uncolored in Phase **I** (1) or not. Next we apply Talagrand's Inequality to prove the following three claims.

- Claim 1: $\Pr[|N(v) \cap R| > \Delta/300] \leq \frac{1}{3\Delta^7}$.
- Claim 2: $\Pr[|N(v) \cap T| > \Delta/300] \leq \frac{1}{3\Delta^7}$.
- Claim 3: $\Pr[|N(v) \cap Q| > \Delta/300] \leq \frac{1}{3\Delta^7}$.

To see that these three claims imply Lemma 2, we observe that if $|N(v) \cap L| > \frac{\Delta}{100}$, then either $|N(v) \cap R| > \frac{\Delta}{300}$, $|N(v) \cap T| > \frac{\Delta}{300}$, or $|N(v) \cap Q| > \frac{\Delta}{300}$.

Proof of Claims 1 and 2: Changing the outcome of each trial affects $|N(v) \cap R|$ by at most 2, and every assignment to trials that results $|N(v) \cap R| \geq k$ can be certified by the outcome of $290k$ trials. Hence Talagrand's Inequality implies that

$$\Pr[|N(v) \cap R| > \frac{\Delta}{1000} + t + 120\sqrt{290\frac{\Delta}{1000}}] \leq 4e^{-1000t^2/8 \times 4 \times 290\Delta}.$$

Since changing the outcome of each trial may add or remove at most two vertices from R , at most 2×290 vertices may be added or removed from T . So it affects $|N(v) \cap T|$ by at most 2×290 . Also every assignment to trials that results $|N(v) \cap T| \geq k$ can be certified by the outcome of $290k$ trials. By Talagrand's Inequality

$$\Pr[|N(v) \cap T| > \frac{\Delta}{1000} + t + 34800\sqrt{290\frac{\Delta}{1000}}] \leq 4e^{-1000t^2/8 \times 4 \times 290^3\Delta}.$$

Substituting $t = \frac{\Delta}{1000}$ in both inequalities completes the proof of Claims 1 and 2.

Proof of Claim 3: To prove this claim, instead of directly applying Talagrand's Inequality to the random variable $|N(v) \cap Q|$, we apply it to the random variable $X_v = |N(v) \setminus Q| = \deg(v) - |N(v) \cap Q|$. Changing the outcome of each trial affects X_v by at most 2, and every outcome of trials that results $X_v \geq k$ can be certified by the outcome of $20k$ trials. Since $X_v = \deg(v) - |N(v) \cap Q|$, and so $\mathbf{E}[X_v] \geq \deg(v) - \frac{\Delta}{1000}$, Talagrand's Inequality implies that

$$\Pr[X_v < \deg(v) - \frac{\Delta}{1000} - t - 120\sqrt{20\Delta}] \leq 4e^{-t^2/8 \times 4 \times 20 \times (\deg(v) - \frac{\Delta}{1000})} \leq 4e^{-t^2/1000\Delta}.$$

Substituting $t = \frac{\Delta}{1000}$ in the inequality above shows that

$$\Pr[X_v < \deg(v) - \frac{\Delta}{300}] \leq \frac{1}{3\Delta^7}.$$

Since $X_v = \deg(v) - |N(v) \cap Q|$, we have

$$\Pr[|N(v) \cap Q| > \frac{\Delta}{300}] \leq \frac{1}{3\Delta^7}.$$

■

In Phase **II**, some of the edges that are incident to vertices in L will be uncolored. Since the other endpoints of these edges may lie in $V(G') - V(H) - L$, to guarantee

that these vertices will remain distinguishable we need to prove a stronger condition than just showing that they are distinguishable in σ_1 . The following lemma is needed to prove Lemma 4 (b) which can imply that with positive probability all vertices in $V(G') - V(H) - L$ will remain distinguishable after applying Phase **II**.

Lemma 3 *For every two adjacent vertices $u, v \in V(G') - V(H)$ where $\deg(u) = \deg(v)$,*

$$\Pr[u \notin L \wedge (|S_{\sigma_1}(u) \triangle S_{\sigma_1}(v)| < 10)] < \frac{1}{\Delta^7}.$$

Proof. It is sufficient to prove that $\Pr[|S_{\sigma_1}(u) \triangle S_{\sigma_1}(v)| < 10 | u \notin L] < \frac{1}{\Delta^7}$. Since we want to prove an upper bound for $\Pr[|S_{\sigma_1}(u) \triangle S_{\sigma_1}(v)| < 10 | u \notin L]$, we can assume that $S_\sigma(u) = S_\sigma(v) = S$. (Remember that σ is the edge coloring before applying Phase **I**.)

Suppose that $\deg_{U_{\sigma_1}}(u) = k$ and $k \geq 20$. If $|S_{\sigma_1}(u) \triangle S_{\sigma_1}(v)| < 10$, then there are at least $k - 10$ colors in $S - S_{\sigma_1}(u)$ which are also in $S - S_{\sigma_1}(v)$. Since there are at most two edges between u and v , the probability of this occurring is at most

$$\binom{k}{k-10} \left(\frac{180}{\Delta}\right)^{k-12} < \frac{1}{\Delta^7}.$$

So

$$\Pr[|S_{\sigma_1}(u) \triangle S_{\sigma_1}(v)| < 10 | u \notin L] < \frac{1}{\Delta^7}.$$

■

Lemma 4 *If we apply Phase **I** to the edge coloring σ , and obtain a partial edge coloring σ_1 , then with positive probability*

- (a): *For every vertex $v \in V(G') - V(H)$, we have $|N(v) \cap L| \leq \frac{\Delta}{100}$,*
- (b): *For every two adjacent vertices $u, v \in V(G') - V(H)$ such that $\deg(u) = \deg(v)$ and $u \notin L$, we have $|S_{\sigma_1}(u) \triangle S_{\sigma_1}(v)| \geq 10$.*

Proof. In order to prove the lemma, we define the following “bad” events

- For a vertex $v \in V(G') - V(H)$, let A_v be the event that $|N(v) \cap L| > \frac{\Delta}{100}$.
- For every edge uv such that $u, v \in V(G') - V(H)$, $\deg(u) = \deg(v)$, let A_{uv} be the event that $u \notin L$ and $|S_{\sigma_1}(u) \triangle S_{\sigma_1}(v)| < 10$.

It is easy to see that each event A_X is mutually independent of all events A_Y such that all vertices in X are in a distance of at least 6 from all vertices in Y . Hence each event is independent of all events but at most $2\Delta^6$ events, and by Lemmas 2

and 3 each event occurs with a probability of at most $\frac{1}{\Delta^7}$. So the Local Lemma implies this lemma. $\blacksquare$

Phase II

1. For every vertex $u \in L$ choose 5 edges uv_i ($1 \leq i \leq 5$) such that $v_i \in V(G') - L$, and uv_i is not an unused edge uniformly at random and uncolor them.

Call this new partial edge coloring σ_2 . For every vertex $u \in V(G') - V(H)$, let UC'_u denote the set of the edges that are incident to u and are uncolored in Phase II.

Lemma 5 *Suppose that σ_1 is a partial edge coloring of G' which satisfies Properties (a) and (b) in Lemma 4. If we apply Phase II to σ_1 , and obtain a partial coloring σ_2 , then with positive probability we have*

- (a): *For every vertex $v \in V(G') - V(H) - L$, $|UC'_v| \leq 4$.*
- (b): *For every two adjacent vertices $u, v \in V(G') - V(H)$ such that $\deg(u) = \deg(v)$, we have $S_{\sigma_2}(u) \neq S_{\sigma_2}(v)$.*

Proof. Suppose that $u, v \in V(G') - V(H)$ are two adjacent vertices such that $\deg(u) = \deg(v)$ and $u \notin L$. Then since by Property (b) in Lemma 4 we know that $|S_{\sigma_1}(u) \triangle S_{\sigma_1}(v)| \geq 10$, (a) implies that $S_{\sigma_2}(u) \neq S_{\sigma_2}(v)$. It is trivial that $|UC'_u| = 5$ for all vertices $u \in L$. So it is sufficient to prove (b') instead of (b), where

- (b'): *For every two adjacent vertices $u, v \in L$ such that $\deg(u) = \deg(v)$, we have $S_{\sigma_2}(u) \neq S_{\sigma_2}(v)$.*

We will apply the Local Lemma to show that with positive probability (a) and (b') hold. We define the following two types of “bad” events.

- For every vertex $u \in V(G') - V(H) - L$, and every five edges $uv_1, uv_2, \dots, uv_5$ such that $v_i \in L$ and uv_i is not an unused edge in σ_1 , let $A_{u, \{v_1, \dots, v_5\}}$ denote the event that $uv_i \in UC'_u$ for all $1 \leq i \leq 5$. Note that since G' is a multigraph, $\{v_1, \dots, v_5\}$ is a multiset and may have duplicate elements.
- For every two adjacent vertices $u, v \in L$ where $\deg(u) = \deg(v)$, let A_{uv} denote the event that $S_{\sigma_2}(u) = S_{\sigma_2}(v)$.

First we should give estimates of $\mathbf{Pr}[A_{u, \{v_1, \dots, v_5\}}]$ and $\mathbf{Pr}[A_{uv}]$. For every vertex $v \in V(G') - V(H)$, we have $\deg(v) \geq \frac{\Delta}{3}$, $|N(v) \cap L| \leq \frac{\Delta}{100}$, and the unused degree of v is at most 290. So every edge uv is uncolored in Phase II with a probability of at most $\frac{5}{(\frac{\Delta}{3} - \frac{\Delta}{100} - 290)} < \frac{20}{\Delta}$, and also two parallel edges are uncolored with a probability of at most $\frac{5}{(\frac{\Delta}{4})} \times \frac{4}{(\frac{\Delta}{4} - 1)}$ which is less than $(\frac{20}{\Delta})^2$. Since $u \notin L$, this implies that

$$\Pr[A_{u,\{v_1,\dots,v_5\}}] \leq \left(\frac{20}{\Delta}\right)^5. \quad (2)$$

Suppose that $S_{\sigma_2}(u) = S_{\sigma_2}(v)$ for two adjacent vertices $u, v \in L$ where $\deg(u) = \deg(v)$. Then since for each of u and v , five new incident edges are uncolored, we have $\deg_{U_{\sigma_1}}(u) = \deg_{U_{\sigma_1}}(v)$. Since $S_{\sigma_2}(u) = S_{\sigma_2}(v)$, all the five edges in UC'_v are determined by the five edges in UC'_u , and since the edges in UC'_v are chosen independently from the edges in UC'_u , we have

$$\Pr[A_{uv}] \leq \frac{1}{\left(\frac{\Delta}{5}\right)} \leq \left(\frac{20}{\Delta}\right)^5. \quad (3)$$

Construct a graph D whose vertices are all the events of the above two types, in which two vertices $A_{u,\{v_1,\dots,v_5\}}$ and $A_{u',\{v'_1,\dots,v'_5\}}$ are adjacent if and only if $\{v_1,\dots,v_5\} \cap \{v'_1,\dots,v'_5\} \neq \emptyset$ (even if $u = u'$), two vertices $A_{u,\{v_1,\dots,v_5\}}$ and A_{vw} are adjacent if and only if $\{v_1,\dots,v_5\} \cap \{v,w\} \neq \emptyset$, and two vertices A_{vw} and $A_{v'w'}$ are adjacent if $\{v,w\} \cap \{v',w'\} \neq \emptyset$. It is easy to see that each vertex of D is mutually independent of all the vertices that are not adjacent to it. Since for every vertex $w \in L$, there are at most $\Delta \binom{\frac{\Delta}{100}}{4}$ events of the form $A_{u,\{w,v_2,\dots,v_5\}}$, and there are at most $\frac{\Delta}{100}$ events of the form A_{vw} , the maximum degree of D is at most

$$5 \left(\Delta \binom{\frac{\Delta}{100}}{4} + \frac{\Delta}{100} \right) < 6\Delta \binom{\frac{\Delta}{100}}{4} < \frac{6\Delta^5}{10^8}.$$

Then since $4 \left(\frac{20}{\Delta}\right)^5 \frac{6\Delta^5}{10^8} < 1$, the Local Lemma implies Lemma 5. $\blacksquare$

Let σ_1 and σ_2 be the partial colorings that are guaranteed to exist by Lemmas 4 and 5. The maximum unused degree of σ_1 is 290. So by Lemma 5 (a) the maximum unused degree of σ_2 is 294. By Vizing's Theorem we can color the unused graph U_{σ_2} with 296 new colors, and obtain a $(\Delta + 298)$ -edge coloring c of G' . By Lemma 5 (b) for every two adjacent vertices $u, v \in V(G') - V(H)$, we have $S_c(u) \neq S_c(v)$.

2.3 Third Step

In this step we begin with the edge coloring $c_1 = c$, and repeatedly modify it to eventually obtain a $(\Delta + 300)$ -*avd*-coloring, c_{final} of G' . We only recolor the edges in H , and since the degree of every vertex in $V(G') - V(H)$ is more than the degree of every vertex in H , all of the vertices in $V(G') - V(H)$ will remain distinguishable.

Obtain c_{k+1} from c_k as in the following. Suppose that for an edge $uv \in H$, $S_{c_k}(u) = S_{c_k}(v)$. Since H does not have any isolated edge, we can assume that $\deg_H(u) \geq 2$. Let $v_1, v_2, \dots, v_r$ be the neighbors of u in H , where $r = \deg_H(u)$. Uncolor all edges $uv_1, uv_2, \dots, uv_r$ to obtain a partial edge coloring c'_k .

Lemma 6 *There exist sets $L(uv_i) \subseteq \{1, \dots, \Delta + 300\}$ for $1 \leq i \leq r$, all of size $r + 300$ such that*

- (a): *In the partial edge coloring c'_k all the colors in $L(uv_i)$ are available for the edge uv_i .*
- (b): *If $v' \neq u$ is adjacent to v_i and $S_{c'_k}(v') \setminus S_{c'_k}(v_i)$ has only one element x , then $x \notin L(uv_i)$.*

Proof. Since $\deg(u), \deg(v_i) \leq \frac{\Delta}{3}$, there are at least $\frac{\Delta}{3} + r + 300$ available colors for uv_i . Furthermore there are at most $\frac{\Delta}{3}$ colors for uv_i which dissatisfy Condition (b). This implies that the desired sets exist. $\blacksquare$

Consider all *proper* completions of c'_k such that the color of each uv_i is chosen from $L(uv_i)$. Note that the number of these completions is at least $(r + 300)(r + 299) \dots 301$. Pick one of them randomly and uniformly, and call it $\hat{c}_k$. Since H does not have any multiple edges, for every v_i , there are at most $(r + 300)(r - 1)!$ different possible choices of $\hat{c}_k$ for which $S_{\hat{c}_k}(u) = S_{\hat{c}_k}(v_i)$. So we have

$$\Pr\left[\bigvee_{1 \leq i \leq r} S_{\hat{c}_k}(u) = S_{\hat{c}_k}(v_i)\right] \leq \sum_{i=1}^r \Pr[S_{\hat{c}_k}(u) = S_{\hat{c}_k}(v_i)] \leq \sum_{i=1}^r \frac{(r - 1)!}{(r + 299) \dots 301} < 1.$$

This implies that there exists a completion c_{k+1} of c'_k such that u is distinguishable, and by Lemma 6 (b) all neighbors of u are also distinguishable. Since we only changed the color of the edges incident to u , the number of indistinguishable vertices is decreased. Hence by repeatedly applying this procedure we will eventually obtain a $(\Delta + 300)$ -avd-coloring of G' .

Acknowledgements

I want to thank my supervisor, Michael Molloy, for reading the draft version of this paper and his valuable comments and suggestions.

References

- [1] S. Akbari, H. Bidkhori, and N. Nosrati. r -strong edge colorings of graphs. *Discrete Math.*, to appear.
- [2] P.N. Balister, E. Györi, J. Lehel, and R.H. Schelp. Adjacent vertex distinguishing edge-colorings. *J. Graph Theory*, to appear.
- [3] P.N. Balister, O.M. Riordan, and R.H. Schelp. Vertex-distinguishing edge colorings of graphs. *J. Graph Theory*, 42:95–109, 2003.

- [4] A. C. Burris and R. H. Schelp. Vertex-distinguishing proper edge-colorings. *J. Graph Theory*, 26 (2):73–82, 1997.
- [5] O. Favaron, H. Li, and R. H. Schelp. Strong edge coloring of graphs,. *Discrete Math.*, 159:103–109, 1996.
- [6] M. Ghandehari and H. Hatami. Two upper bounds for the strong edge chromatic number. *Preprint*.
- [7] M. Horňák and R. Soták. Asymptotic behavior of the observability of Q_n . *Discrete Math.*, 176:139–148, 1997.
- [8] M. Karoński, T. Łuczak, and A. Thomason. Edge weights and vertex colours. *J. Combin. Theory Ser. B*, 91:151–157, 2004.
- [9] M. Molloy and B. Reed. *Graph Colouring and the Probabilistic Method*. Springer, 2002.
- [10] D. B. West. *Introduction to Graph Theory*. Prentice-Hall, Inc, United States of American, 2001. 2nd Edition.
- [11] Z. Zhang, L. Liu, and J. Wang. Adjacent strong edge coloring of graphs. *Appl. Math. Letters*, 15:623–626, 2002.